

Models of PT^- with internal induction for total formulae

Cezary Cieśliński* Mateusz Łelyk† Bartosz Wcisło‡

Institute of Philosophy, University of Warsaw, Poland

Abstract

We show that a typed compositional theory of positive truth with internal induction for total formulae (denoted by PT_{tot}) is not semantically conservative over Peano arithmetic. In addition, we observe that a class of models of PA expandable to models of PT_{tot} contains every recursively saturated model of arithmetic. Our results point to a gap in the philosophical project of describing the use of the truth predicate in model-theoretic contexts.

For quite a while conservativity has been promoted as an important trait of deflationary truth theories. To our knowledge, the idea was introduced for the first time by Leon Horsten, who declared conservativity to be a commitment of Horwich’s ‘minimal theory’.¹ Since then Horsten’s proposal of explaining the deflationary tenet of ‘thinness’ or ‘neutrality’ of truth in terms of conservativity has been often repeated, discussed and refined in logical and philosophical literature.

At least two notions of conservativity have been proposed as tools for the deflationist to explicate his position. One notion is syntactic; the intended meaning is that a conservative extension does not prove new theorems of the base language. The second is semantic and concerns the possibility of expanding models.²

*c.cieslinski@uw.edu.pl

†mlelyk@student.uw.edu.pl

‡bar.wcislo@gmail.com

¹In Horsten’s own words: ‘The minimalist theory entails that a truth predicate should be conservative over a given theory that is stated without the truth predicate (or any other semantical notions).’ See [12], p. 183.

²See [3] for a discussion of the philosophical motivation behind both versions of the conservativity demand for deflationary theories of truth.

Definition 1. Let T_1 and T_2 be theories in languages L_1 and L_2 (with $L_1 \subseteq L_2$). Then:

- (i) T_2 is syntactically conservative over T_1 iff $\forall \psi \in L_1 [T_2 \vdash \psi \rightarrow T_1 \vdash \psi]$.
- (ii) T_2 is semantically conservative over T_1 iff every model of T_1 can be expanded to a model of T_2 .

Although semantic conservativity entails conservativity in the syntactic sense, these two notions are not equivalent. There are well-known examples of axiomatic truth theories which are syntactically, but not semantically conservative over their base theories of syntax.

Research on axiomatic theories of truth permits us to recognise the following fairly general patterns, exemplified by many truth theories.

- Axiomatic truth theories which are both compositional and fully inductive are not syntactically conservative over their base theories of syntax. However, the lack of even one of these properties typically produces syntactic conservativity.³
- Even without compositionality, full induction in the language with the truth predicate typically produces semantic non-conservativity of an axiomatic theory of truth. However, it is known that compositionality by itself (that is, without extended induction) can be squared with semantic conservativity.⁴

The initial reception of these and related results has been mainly negative. Some authors (see in particular [16] and [14]) employed the conservativity demand as a weapon against deflationary theories of truth, arguing that deflationary truth *should* be conservative, but immediately adding that *it cannot be*, because conservative truth theories are too weak. However, more recently some defences of conservative truth theories - even in a stronger semantic sense of the word - have been put forward in the literature. In particular, in [8] Fischer and Horsten proposed to study axiomatic truth theories treated as characterisations of the use of the truth predicate in model-theoretic contexts. In their own words:

³Compositional theories CT, KF and FS are syntactically non-conservative over PA, but they become conservative as soon as the extended induction - that is, induction for formulae with the truth predicate - is removed (see [11] for further details). On the other hand, disquotational non-compositional axiomatic theories TB, UTB and PTB are syntactically conservative, even though they contain full extended induction (see [2] and [4]). A curious counterexample is a disquotational theory PUTB discussed in [10], which is not syntactically conservative over Peano arithmetic.

⁴If a theory of truth proves biconditionals ' $T(\varphi) \equiv \varphi$ ' for all arithmetical sentences φ and in addition contains full extended induction, then it is not semantically conservative over PA (see [2]). On the other hand, compositional positive truth axioms without any extended induction produce theories which are semantically conservative over PA (see [11] for the details).

There are contexts where one is reluctant to privilege one model over another, and where one does not want a theory of truth to exclude models for the original language. In particular, this is the case in the single mathematical field where truth predicates play a major role, viz. *model theory*, and in uses of model theory in proof theory. ([8], p. 345)

In effect, Fischer and Horsten describe their endeavour as similar in important respects to that of Tarski, who attempted to establish ‘beyond reasonable doubt that the uses of truth predicates in metamathematics are legitimate.’ ([8], p. 345)

Apart from semantic conservativity, there is also another desirable trait of the aforementioned ‘uses of truth predicates’. Truth has expressive power - it ‘widens the class of thoughts that we can express’ ([8], p. 345). One indication of the expressive power is the non-interpretability of our theory of truth in its base theory of syntax; there is then a precise sense in which truth brings conceptually something new. Another indication is the non-elementary speed-up of the theory of truth with respect to its arithmetical base theory. Here the expressive power of truth manifests itself in the instrumental value of the truth predicate, namely, truth permits us to shorten proofs. All in all, the moral is that we should search for an axiomatic theory of truth Th which jointly satisfies the following requirements:

- (a) Th is semantically conservative over its arithmetical base theory of syntax B ,
- (b) Th is not interpretable in B ,
- (c) Th has non-elementary speed-up over B .

Do we have at our disposal an axiomatic truth theory which would capture the uses of the truth predicate in model theory? It has been suggested that a theory of typed positive truth with internal induction for total formulae (denoted as PT_{tot} in this paper⁵) fills the bill. Below we introduce the relevant definitions.

Let L_{PA} be the language of Peano arithmetic, with Var and Tm^c being (respectively) the sets of variables and constant arithmetical terms. By L_T we denote the extension of L_{PA} with the new one-place predicate ‘ $T(x)$ ’. For the sentences of L_{PA} the notation ‘ $Sent_{L_{PA}}$ ’ will be used. By \underline{x} we mean the x -th numeral, i.e. the only numeral denoting the number x . We define:

⁵Some authors, notably Fischer in [6], have been using the notation “PT[−]” instead of our PT_{tot} . We prefer a different notation, reserving “ Th ” for truth theories Th without any induction for formulae containing the truth predicate.

Definition 2. PT^- is the theory in the language L_T which is axiomatised by the usual axioms of Peano arithmetic (PA), together with the following truth theoretic axioms:

- (1) $\forall s \forall t \in Tm^c (T(s = t) \equiv val(s) = val(t))$
- (2) $\forall s \forall t \in Tm^c (T(\neg s = t) \equiv val(s) \neq val(t))$
- (3) $\forall \psi \in Sent_{L_{PA}} (T(\neg \neg \psi) \equiv T(\psi))$
- (4) $\forall \varphi \forall \psi \in Sent_{L_{PA}} (T(\varphi \wedge \psi) \equiv T(\varphi) \wedge T(\psi))$
- (5) $\forall \varphi \forall \psi \in Sent_{L_{PA}} (T(\neg(\varphi \wedge \psi)) \equiv T(\neg \varphi) \vee T(\neg \psi))$
- (6) $\forall v \in Var \forall \varphi \in L_{PA} (T(\forall v \varphi) \equiv \forall x T(\varphi(\underline{x}/v)))$
- (7) $\forall v \in Var \forall \varphi \in L_{PA} (T(\neg \forall v \varphi) \equiv \exists x T(\neg \varphi(\underline{x}/v)))$

The above axiomatisation is the same as presented in [6]. In [8] the considered theory has been augmented with two new axioms to the effect that truth is extensional and only sentences are true, i.e with the following **extensionality principle**

$$\forall \phi \in L_{PA} \forall s, t \in Tm^c (val(t) = val(s) \rightarrow (T(\phi(t)) \equiv T(\phi(s)))) \quad (\text{EXT})$$

and the **normality principle**

$$\forall x (T(x) \rightarrow x \in Sent_{L_{PA}}). \quad (\text{NORM})$$

Although at some point we will consider also these additional axioms, we emphasise that in the terminology adopted in this paper they do not belong to PT^- proper.

In the next move we extend PT^- with a weak form of induction for total arithmetical formulae.

Definition 3. Let $\text{tot}(\phi)$ be a shorthand for $\forall x [T(\phi(x)) \vee T(\neg \phi(x))]$. By the principle of internal induction for total formulae (Ind_{tot}) we mean the following single sentence of the language L_T :

$$\forall \phi(x) \in L_{PA} \left(\text{tot}(\phi(x)) \rightarrow \left(\forall x [T(\phi(\underline{x})) \rightarrow T(\phi(\underline{x}+1))] \rightarrow (T(\phi(\underline{0})) \rightarrow \forall x T(\phi(\underline{x}))) \right) \right),$$

where the quantifier ' $\forall \phi(x)$ ' reads 'for every formula $\phi(x)$ with at most x free'. We read the expression ' $\text{tot}(x)$ ' as ' x is **total**'. We denote the theory $PT^- + Ind_{tot}$ by PT_{tot} .

One could wonder whether such a form of induction is not overly restrictive. Why should we have induction for total formulae only? However, Fischer and Horsten claim that the restriction becomes natural and well-motivated as soon as we appreciate that it is *all* models of arithmetic - including the nonstandard ones - that matter:

We do not want to accept instantiations of induction for non-standard elements that are not truth-determinate for the property in question for exactly the same reason that we resist inductive premises for soritical predicates. Note that the reply that in the “intended” model there are no such non-standard elements to be found is not undermining our motivation for restriction; as we have emphasised repeatedly, we are adopting the model-theoretic viewpoint, and from this viewpoint, all models are on a par. ([8], p. 355)

In [6] it has been claimed that PT_{tot} is semantically conservative over PA; in the same paper the author proves that PT_{tot} is not interpretable in its base theory (PA). In turn, in [5] it is shown that PT_{tot} has a non-elementary speed-up over PA. Taken together, these results imply that PT_{tot} is indeed an excellent candidate for the role of the theory characterising the use of the truth predicate in model-theoretic contexts.

Unfortunately, the conservativity proof presented in [6] contains a flaw.⁶ Indeed, the main result of the present paper is that PT_{tot} is not semantically conservative over Peano arithmetic (hence the same is true about PT_{tot} with (EXT) and (NORM) added). In view of this, the question still remains whether we have at our disposal a natural axiomatic truth theory satisfying requirements (a)-(c).

An easy argument based on the existence of fixed points for monotone operators shows that PT^- is model-theoretically conservative over PA (see [11]). However, it transpires that adding internal induction for total formulae comes with a price. The following theorem states the semantic non-conservativity of PT_{tot} .

Theorem 4. *There is a model of PA which cannot be expanded to a model of PT_{tot} .*

Before we proceed to the proof, we shall define one construction in propositional logic which is very useful in the context of investigating compositional theories of truth with restricted induction. The construction has originally appeared in [17] although its properties were not spelled out in full generality.

⁶We are grateful to Martin Fischer for the email correspondence concerning this matter.

Definition 5. Let $\alpha = (\alpha_0 \dots \alpha_c)$ and $\beta = (\beta_0 \dots \beta_c)$ be arbitrary sequences of formulae. By a **disjunction of β_i** (for $0 \leq k \leq n \leq c$) **with stopping condition α** , denoted

$$\bigvee_{i=k}^{n,\alpha} \beta_i$$

we mean any of the following formulae defined by backward induction on k :

1. $\bigvee_{i=n}^{n,\alpha} \beta_i = (\alpha_n \wedge \beta_n).$
2. $\bigvee_{i=k}^{n,\alpha} \beta_i = \neg(\alpha_k \wedge \neg\beta_k) \wedge \left((\alpha_k \wedge \beta_k) \vee \bigvee_{i=k+1}^{n,\alpha} \beta_i \right).$

The intuition behind the above formulae is as follows: we want to search through i as long as we do not see i_0 such that α_{i_0} is true. Then we stop and check whether β_{i_0} is satisfied. If it is, then the whole formula is true, if not, then the whole formula is false, regardless of the truth value of β_j for $j > i_0$. We call it a disjunction, since if α is chosen so that exactly one of the α_i -s is true, then the above construction is equivalent in propositional logic to the disjunction:

$$\bigvee_{i=k}^n (\alpha_i \wedge \beta_i).$$

Although one should note that in this case it is also equivalent to

$$\bigwedge_{i=k}^n (\alpha_i \rightarrow \beta_i).$$

So, in a sense, it is a propositional analogue of a Δ_1 -formula.

Given a model M of Peano arithmetic and sequences $\alpha, \beta \in M$,⁷ the construction can be reproduced inside M . In such a case the expression ' $\bigvee_{i=k}^{n,\alpha} \beta_i$ ' will refer to the unique, possibly nonstandard formula in the sense of M . The most important property of the above construction, one we have already alluded to, is that formulae obtained in this way (even nonstandard ones!) behave well in models of compositional truth theories without induction for the language with the truth predicate. These nice properties are encapsulated in the following lemma:

⁷More exactly, we assume here that α and β are elements of M such that for some $c \in M$, the formal analogue of the statement ' α and β are sequences of length c containing arithmetical formulae' is true in M .

Lemma 6. *Let M be an arbitrary nonstandard model of PT^- . Let $\alpha, \beta \in M$ be sequences of nonstandard length c , containing as elements arithmetical formulae $\alpha_i(x), \beta_i(y)$ with the free variables indicated.⁸ Suppose that for all $i \in \omega$ the formulae $\alpha_i(x)$ are standard. Let $a \in M$ be such that $j_0 \in \omega$ is the smallest natural number satisfying the condition $M \models T(\alpha_{j_0}(a))$. Then for every $k \leq j_0$ and for every $b \in M$ the following holds:*

$$(a) \quad M \models T\left(\bigvee_{i=k}^{c,\alpha} \beta_i(a, b)\right) \equiv T(\beta_{j_0}(b)).$$

$$(b) \quad M \models T\left(\neg \bigvee_{i=k}^{c,\alpha} \beta_i(a, b)\right) \equiv T(\neg \beta_{j_0}(b)).$$

In the proof we will use the well-known fact that for every standard formula $\varphi(x_1 \dots x_n)$:

$$\text{PT}^- \vdash \forall x_1 \dots x_n \left(T(\varphi(x_1 \dots x_n)) \equiv \varphi(x_1 \dots x_n) \right).$$

It immediately follows that for every standard formula $\varphi(x_1 \dots x_n)$:

$$\text{PT}^- \vdash \forall x_1 \dots x_n \left(T(\neg \varphi(x_1 \dots x_n)) \equiv \neg T(\varphi(x_1 \dots x_n)) \right).⁹$$

Proof. We prove the lemma by backward metainduction on k . Starting with $k = j_0$, note that there exists γ such that the disjunction of β_i with stopping condition α from j_0 to c may be written as:

$$\bigvee_{i=j_0}^{c,\alpha} \beta_i(a, b) = \underbrace{\neg(\alpha_{j_0}(a) \wedge \neg \beta_{j_0}(b))}_A \wedge \underbrace{((\alpha_{j_0}(a) \wedge \beta_{j_0}(b)) \vee \gamma)}_B.$$

For the implication from right to left in part (a) of the lemma, assume that $M \models T(\beta_{j_0}(b))$. Therefore by the axiom of PT^- for double negation, we obtain $M \models T(\neg \neg \beta_{j_0}(b))$, which in turn (by compositional axiom for negated conjunction) permits us to conclude that $M \models T(A)$. Since by assumption $M \models T(\alpha_{j_0}(a))$, we obtain also $M \models T(\alpha_{j_0}(a) \wedge \beta_{j_0}(b))$. Applying again compositional axioms of PT^- we get $M \models T(A \wedge B)$, as required.

For the implication from right to left in (b) it is enough to observe that $M \models T(\neg \beta_{j_0}(b))$ together with $M \models T(\alpha_{j_0}(a))$ gives us immediately $M \models T(\neg A)$, and so $M \models T(\neg(A \wedge B))$.

Proving the opposite implication in (a), assume that $M \models T(A \wedge B)$. From $M \models T(A)$ by compositional axioms of PT^- we obtain: $M \models T(\neg \alpha_{j_0}(a))$ or

⁸Again, this is taken to mean that all elements of α and β are formulae in the sense of M .

⁹In effect, we easily obtain the information that for every standard formula $\varphi(x)$, truth is provably total and consistent in PT^- . In other words, if $\varphi(x)$ is standard, then $\text{PT}^- \vdash \text{tot}(\varphi(x)) \wedge \neg \exists x (T(\varphi(x)) \wedge T(\neg \varphi(x)))$.

$M \models T(\neg\neg\beta_{j_0}(b))$. But $M \models T(\alpha_{j_0}(a))$, and since α_{j_0} is standard, this gives us $M \models \neg T(\neg\alpha_{j_0}(a))$, and so by the compositional axiom for double negation, $M \models T(\beta_{j_0}(b))$.

Proving the opposite implication in (b), assume that $M \models T(\neg(A \wedge B))$, so $M \models T(\neg A) \vee T(\neg B)$. If $M \models T(\neg A)$, then by compositional axioms $M \models T(\neg\beta_{j_0}(b))$. If $M \models T(\neg B)$, then by compositional axioms $M \models T(\neg\alpha_{j_0}(a)) \vee T(\neg\beta_{j_0}(b))$. But $M \models T(\alpha_{j_0}(a))$, so (since α_{j_0} is standard) $M \models \neg T(\neg\alpha_{j_0}(a))$ and therefore $M \models T(\neg\beta_{j_0}(b))$.

Suppose now that our claim is true for a given $k+1 \leq j_0$. In other words, we have:

- (i) $M \models T\left(\bigvee_{i=k+1}^{c,\alpha} \beta_i(a, b)\right) \equiv T(\beta_{j_0}(b))$.
- (ii) $M \models T\left(\neg \bigvee_{i=k+1}^{c,\alpha} \beta_i(a, b)\right) \equiv T(\neg\beta_{j_0}(b))$.

Our task is to prove the claim for k . Observe that by Definition 5:

$$\bigvee_{i=k}^{c,\alpha} \beta_i = \underbrace{\neg(\alpha_k(a) \wedge \neg\beta_k(b))}_C \wedge \underbrace{((\alpha_k(a) \wedge \beta_k(b)) \vee \bigvee_{i=k+1}^{c,\alpha} \beta_i(a, b))}_D.$$

For the implication from right to left in part (a) of the lemma, assume that $M \models T(\beta_{j_0}(b))$. Therefore by (i) we have: $M \models T\left(\bigvee_{i=k+1}^{c,\alpha} \beta_i(a, b)\right)$, so $M \models T(D)$. Since $k < j_0$, by the choice of j_0 we have $M \models \neg T(\alpha_k(a))$; therefore $M \models T(\neg\alpha_k(a))$ because α_k is standard. In effect, the compositional axiom of PT^- for negated conjunction permit us to conclude also that $M \models T(C)$ and thus $M \models T(C \wedge D)$.

For the implication from right to left in (b), assume that $M \models T(\neg\beta_{j_0}(b))$. Then by (ii) $M \models T\left(\neg \bigvee_{i=k+1}^{c,\alpha} \beta_i(a, b)\right)$. Since $M \models \neg T(\alpha_k(a))$, the compositional axioms of PT^- give us $M \models T(\neg(\alpha_k(a) \wedge \beta_k(b)))$, so $M \models T(\neg D)$ and finally $M \models T(\neg(C \wedge D))$ as required.

For the implication from left to right in (a), assume that $M \models T(C \wedge D)$, so in particular $M \models T(\alpha_k(a) \wedge \beta_k(b)) \vee T\left(\bigvee_{i=k+1}^{c,\alpha} \beta_i(b)\right)$. But $M \models \neg T(\alpha_k(a))$ and therefore the second disjunct must be true, which by (i) permits us to conclude that $M \models T(\beta_{j_0}(b))$.

For the implication from left to right in (b), assume that $M \models T(\neg(C \wedge D))$, so $M \models T(\neg C) \vee T(\neg D)$. However, by compositional axioms $M \models T(\neg C) \equiv (T(\alpha_k(a)) \wedge T(\neg\beta_k(b)))$ and since $M \models \neg T(\alpha_k(a))$, we obtain $M \models T(\neg D)$. Then by the compositional axiom of PT^- for negated disjunction it follows that $M \models T\left(\neg \bigvee_{i=k+1}^{c,\alpha} \beta_i(a, b)\right)$ and therefore by (ii) $M \models T(\neg\beta_{j_0}(b))$.

□

Corollary 7. *Let M , α , β , a and j_0 satisfy the assumptions of Lemma 6. If in addition the formula $\beta_{j_0}(y)$ is standard, then $M \models \forall k \leq j_0 \text{ tot} \left(\bigvee_{i=k}^{c, \alpha} \beta_i(a, y) \right)$.*

Proof. It is enough to observe that if $\beta_{j_0}(y)$ is standard, then $M \models \forall b T \left((\beta_{j_0}(b)) \vee T(\neg \beta_{j_0}(b)) \right)$. Then by Lemma 6 the corollary follows immediately. \square

Now we are ready to prove our theorem. The argument will consist in showing that no nonstandard prime model of Peano arithmetic is expandable to a model of PT_{tot} .¹⁰

Proof of Theorem 4. Let K be an arbitrary nonstandard prime model of Peano arithmetic. From now on we take for granted that all elements of K are definable in K by arithmetical formulae without parameters. The claim will be that K is not expandable to a model of PT_{tot} . Suppose for contradiction that $(K, T) \models \text{PT}_{\text{tot}}$. We will argue that in such a case there is an element e of K which codes $Th(K)$ (that is, e codes the set of all arithmetical sentences true in K). However, such an element e cannot exist in a prime model, because then e would be definable in K , generating a contradiction with Tarski's undefinability theorem.

For starters, fix any nonstandard $c \in K$ and let α be a recursive enumeration of formal definitions up to c . That is, given a fixed recursive enumeration $\phi_0(x), \phi_1(x) \dots$ of arithmetical formulae with exactly one free variable x , each $\alpha_i(x)$ has the form:

$$\phi_i(x) \wedge \forall y < x \neg \phi_i(y).$$

For $a \in K$, we say that $\alpha_i(x)$ defines a in K iff i is the smallest natural number such that $K \models \alpha_i(a)$. Observe that since K is prime, such a number exists for every $a \in K$ and a corresponding formula $\alpha_i(x)$ is standard.

We define now the second sequence β , containing formulae $\beta_i(y)$ with one free variable, characterised as:

$$\forall \phi < i [Sent_{L_{PA}}(\phi) \rightarrow (\phi \in y \equiv T_i(\phi))].$$

The expression ' T_i ' stands for an arithmetical truth predicate for sentences below i . The exact shape of this predicate is not of crucial importance; what

¹⁰Given a model M of Peano arithmetic, the universe of a prime model K can be defined as the set of all those elements of M which are definable in M by arithmetical formulae without parameters. In turn, the operations of K are defined as those of M restricted to the universe of K . For further information about prime models we refer the reader to [13], p. 91ff.

will really matter is that for $i \in \omega$, $T_i(x)$ - and therefore also $\beta_i(y)$ - is a standard arithmetical formula.¹¹

We define:

- $\psi(x, y) := \bigvee_{i=0}^{c, \alpha} \beta_i(x, y)$,
- $\xi(z) := \exists y \forall x < z \psi(x, y)$.

We emphasise that $\psi(x, y)$ and $\xi(z)$ are defined in a model K and not externally. With c being nonstandard, both $\psi(x, y)$ and $\xi(z)$ should be thought of as nonstandard arithmetical formulae - elements of K which are *perceived* by K as formulae, not to be confused with the ‘real world’ arithmetical expressions.¹²

We are going to show that:

- (i) $(K, T) \models \text{tot}(\xi(z))$,
- (ii) $(K, T) \models T(\xi(0)) \wedge \forall x \left(T(\xi(x)) \rightarrow T(\xi(x+1)) \right)$.

For (i), first observe that $(K, T) \models \forall a \text{ tot}(\psi(a, y))$. In order to see this, fix $a \in K$ and let $\alpha_{j_0}(x)$ define a . Fixing b , it is enough to observe that the assumptions of Corollary 7 are satisfied; therefore $(K, T) \models \forall k \leq j_0 \text{ tot}\left(\bigvee_{i=k}^{c, \alpha} \beta_i(a, y)\right)$. For $k = 0$, this gives $(K, T) \models \forall a \text{ tot}(\psi(a, y))$.

It easily follows that $\forall a \in K (K, T) \models \text{tot}(\forall x < a \psi(x, y))$.¹³ Finally, we argue for the totality of $\xi(z)$. Given an arbitrary $a \in K$, if $(K, T) \models \exists y T(\forall x < a \psi(x, y))$, then $(K, T) \models T(\xi(a))$. Otherwise $(K, T) \models \forall y \neg T(\forall x < a \psi(x, y))$, which (by totality of ‘ $\forall x < a \psi(x, y)$ ’) entails $(K, T) \models T(\neg \xi(a))$.

For (ii), obviously $(K, T) \models T(\xi(0))$,¹⁴ so we move to the second conjunct. Fix b such that $(K, T) \models T(\xi(b))$; in other words: $(K, T) \models T(\exists y \forall x < b \psi(x, y))$. Choose $e \in K$ such that $(K, T) \models T(\forall x < b \psi(x, e))$. Our task is to obtain $e' \in K$ such that $(K, T) \models T(\forall x < b+1 \psi(x, e'))$.

Let $\alpha_i(x)$ be the least definition of b . We put:

¹¹For example, predicates $T_i(x)$ could be defined as ‘ $(x = \ulcorner \psi_0 \urcorner \wedge \psi_0) \vee \dots \vee (x = \ulcorner \psi_m \urcorner \wedge \psi_m)$ ’, where $\psi_0 \dots \psi_m$ are all arithmetical sentences with gödel numbers smaller than i . Note that this construction of $T_i(x)$ can be carried out in an arbitrary model M of PA. In effect, we will have formulae (in the sense of M) $T_c(x)$ for an arbitrary element c of M , even a nonstandard one. However, it is important to emphasise that in each formula $T_i(x)$ the letter ‘ i ’ is not a variable. The index gives rather the information that in the corresponding formula all sentences with gödel numbers smaller than (fixed) i are taken into account.

¹²Accordingly, it would make no sense to say, for example, that $\psi(x, y)$ is satisfied in K by some objects a and b . Nevertheless, nonstandard formulae can be employed in the scope of the truth predicate and we are allowed to say, for example, that $(K, T) \models T(\psi(a, b))$.

¹³Fix a and b ; assume that $(K, T) \models \neg T(\forall x < a \psi(x, b))$. Then $(K, T) \models \exists x < a \neg T(\psi(x, b))$. Choosing such an $x \in K$, we obtain $(K, T) \models \neg T(\psi(x, b))$ and by totality of $\psi(a, y)$ for every a , we obtain $(K, T) \models T(\neg \psi(x, b))$ and thus $(K, T) \models T(\neg \forall x < a \psi(x, b))$.

¹⁴It is enough to observe that there is no $x < 0$ in K , so an arbitrary y is a witness for ‘ $\exists y \forall x < 0 \psi(x, y)$ ’.

$$e' = e - \{0 \dots i - 1\} \cup \{\psi \in \text{Sent}_{L_{PA}} \mid \psi < i \wedge K \models \psi\}.$$

All that remains for the completion of the proof of (ii) is to show that for every $d < b + 1$, $(K, T) \models T(\psi(d, e'))$. Fix such a d and let $\alpha_j(x)$ be the least definition of d . By Lemma 6, it is enough to obtain $(K, T) \models T(\beta_j(e'))$; in other words, we want to have:

$$(*) \quad (K, T) \models T\left(\forall \phi < j [\text{Sent}_{L_{PA}}(\phi) \rightarrow (\phi \in e' \equiv T_j(\phi))]\right).$$

Since $d < b + 1$, we consider two cases. If $d = b$, then $j = i$ and $(*)$ follows trivially from the definition of e' . On the other hand, if $d < b$, then $(K, T) \models T(\psi(d, e))$, and so by Lemma 6 $(K, T) \models T(\beta_j(e))$. In other words, we have:

$$(**) \quad (K, T) \models T\left(\forall \phi < j [\text{Sent}_{L_{PA}}(\phi) \rightarrow (\phi \in e \equiv T_j(\phi))]\right).$$

It is easy now to show that $(*)$ must hold. Fixing $\phi < j$, we observe that if $\phi < i$, then $(K, T) \models T(\phi \in e' \equiv T_j(\phi))$ because by definition, e' codes only true sentences below i . Otherwise $\phi \geq i$, but then $\phi \in e$ and $\phi \in e'$, so by $(**)$ we also conclude that $(K, T) \models T(\phi \in e' \equiv T_j(\phi))$.

This finishes the proofs of (i) and (ii). At this point we know that $\xi(z)$ is total and inductive in (K, T) , so by the axiom of internal induction we conclude that $(K, T) \models \forall z T(\xi(z))$. Let a be a nonstandard element of K . Since $(K, T) \models T(\xi(a))$, we can choose e such that $(K, T) \models \forall x < a T(\psi(x, e))$.

We will show that e codes $Th(K)$ in K ; in other words, we show that:

$$(***) \quad \forall n \in \omega \quad K \models \forall \phi < n [\text{Sent}_{L_{PA}}(\phi) \rightarrow (\phi \in e \equiv T_n(\phi))].$$

Fix $n \in \omega$. Let k and i be elements of ω such that $\alpha_i(x)$ defines k in K and $i > n$. Since $k < a$,¹⁵ we have: $(K, T) \models T(\psi(k, e))$, which by Lemma 6 is equivalent to $(K, T) \models T(\beta_i(e))$. In other words:

$$(K, T) \models T\left(\forall \phi < i [\text{Sent}_{L_{PA}}(\phi) \rightarrow (\phi \in e \equiv T_i(\phi))]\right).$$

Applying disquotation (valid in PT^- for standard formulae with parameters), we obtain:

$$K \models \forall \phi < i [\text{Sent}_{L_{PA}}(\phi) \rightarrow (\phi \in e \equiv T_i(\phi))].$$

Since $i > n$, $(***)$ follows trivially. In effect, e codes $Th(K)$ in K . But this is impossible in prime models, thus a contradiction is obtained and the proof is finished. \square

¹⁵We remind that k is standard and a is a nonstandard element of K .

In this section we further approximate the class of those models of PA that admit an expansion to models of PT_{tot} . In general such a class is a handy tool for comparing properties of axiomatic theories of truth. Let us introduce the precise definition and a piece of notation:

Definition 8. Let Th be any extension of PA (possibly in extended language). By \mathfrak{Th} we denote the class of models of PA that can be expanded to a model of Th .

For example, it is known that for every nonstandard model \mathcal{M} , $\mathcal{M} \in \mathfrak{TB}$ if and only if \mathcal{M} codes its own theory, i.e.

$$\{\ulcorner \phi \urcorner \mid \phi \in \text{Sent}_{L_{PA}} \wedge \mathcal{M} \models \phi\}$$

is coded in \mathcal{M} .¹⁶

Results on such classes can be used to obtain some information about *relative truth definability*¹⁷ between axiomatic theories of truth. The connection between these two notions is established via the following fact:

Fact 9. Let Th_1, Th_2 be two axiomatic truth theories. If Th_1 is relatively truth definable in Th_2 , then $\mathfrak{Th}_2 \subseteq \mathfrak{Th}_1$.

For the (immediate) proof, see [9].

In the previous section we showed that the class of prime models of PA is disjoint from $\mathfrak{PT}_{\text{tot}}$. In the next theorem we approximate this class from below (by \mathfrak{RS} we denote the class of recursively saturated models of PA). For technical simplicity, we will prove the theorem for the stronger theory $\text{PT}_{\text{tot}} + (\text{EXT}) + (\text{NORM})$, which from now on we will denote by RPT_{tot} (**regular** PT_{tot}).

Theorem 10. $\mathfrak{RS} \subseteq \mathfrak{RPT}_{\text{tot}}$.¹⁸

Note that since PT_{tot} is a subtheory of RPT_{tot} , the above result clearly implies that every recursively saturated model of PA can be expanded to a model of PT_{tot} .

¹⁶For the details, see [2]. However, it should be noted that our ability to provide a purely arithmetical characterisation of \mathfrak{TB} makes TB rather exceptional. For most axiomatic truth theories we are only able to restrict the class of possible candidates. For example, it is known that the class of recursively saturated models of PA properly contains \mathfrak{CT}^- and \mathfrak{UTB} .

¹⁷For the definition of this notion and the discussion of its philosophical relevance, see [9].

¹⁸Cf. [1], where it is argued that every recursively saturated model is expandable to a model of an untyped truth theory KF_t . Cantini's KF_t is formulated in the language with two predicates ' T ' and ' F ' (for truth and falsity respectively); it contains also the axiom of internal induction for total formulae.

The proof follows immediately from two lemmata, which we consider interesting also for their own sake. Before stating them let us introduce two more notions (the expressions $Sent_{\mathcal{M}}$, $Tm_{\mathcal{M}}^c$, $Form_{\mathcal{M}}^1$ denote the set of arithmetical sentences, closed terms and arithmetical formulae with at most one free variable in the sense of model \mathcal{M}):

Definition 11. By the **term formulation** of PT^- , denoted by tPT^- , we mean PT^- with the quantifier axioms:

$$(6) \quad \forall v \in Var \forall \varphi \in L_{PA} (T(\forall v \varphi) \equiv \forall x T(\varphi(\underline{x}/v)))$$

$$(7) \quad \forall v \in Var \forall \varphi \in L_{PA} (T(\neg \forall v \varphi) \equiv \exists x T(\neg \varphi(\underline{x}/v)))$$

replaced with the following ones:

$$(6') \quad \forall v \in Var \forall \varphi \in L_{PA} (T(\forall v \varphi) \equiv \forall t \in Tm^c T(\varphi(t/v)))$$

$$(7') \quad \forall v \in Var \forall \varphi \in L_{PA} (T(\neg \forall v \varphi) \equiv \exists t \in Tm^c T(\neg \varphi(t/v))).$$

We define the term formulation of PT_{tot} as the term formulation of PT^- extended with the internal induction axiom (Ind_{tot}). We note that there is no need to define the term formulations of RPT or RPT_{tot} , since over any extension of PT^- containing (EXT) the conditions 6 (7) and 6' (7') are equivalent.

Definition 12. Let $\mathcal{M} \models PA$, let $A \subseteq M$ and let ϕ be a L_{PA} sentence in the sense of \mathcal{M} . We define:

$$\begin{aligned} \Theta_{\mathcal{M}}(\phi, A) := & \mathcal{M} \models \exists s, t \in Tm^c [\phi = (s = t) \wedge val(s) = val(t)] \\ & \vee \mathcal{M} \models \exists s, t \in Tm^c [\phi = \neg(s = t) \wedge val(s) \neq val(t)] \\ & \vee \exists \psi \in Sent_{\mathcal{M}} [\mathcal{M} \models \phi = \neg \neg \psi] \wedge \psi \in A \\ & \vee \exists \psi_1, \psi_2 \in Sent_{\mathcal{M}} [\mathcal{M} \models \phi = (\psi_1 \wedge \psi_2)] \wedge (\psi_1 \in A \wedge \psi_2 \in A) \\ & \vee \exists \psi_1, \psi_2 \in Sent_{\mathcal{M}} [\mathcal{M} \models \phi = \neg(\psi_1 \wedge \psi_2)] \wedge (\neg \psi_1 \in A) \vee (\neg \psi_2 \in A) \\ & \vee \exists \psi_1, \psi_2 \in Sent_{\mathcal{M}} [\mathcal{M} \models \phi = (\psi_1 \vee \psi_2)] \wedge (\psi_1 \in A) \vee (\psi_2 \in A) \\ & \vee \exists \psi_1, \psi_2 \in Sent_{\mathcal{M}} [\mathcal{M} \models \phi = \neg(\psi_1 \vee \psi_2)] \wedge (\neg \psi_1 \in A) \wedge (\neg \psi_2 \in A) \\ & \vee \exists \psi(x) \in Form_{\mathcal{M}}^1 [\mathcal{M} \models \phi = \exists x \psi] \wedge \exists s \in Tm^c (\psi(s) \in A) \\ & \vee \exists \psi(x) \in Form_{\mathcal{M}}^1 [\mathcal{M} \models \phi = \neg \exists x \psi] \wedge \forall s \in Tm^c (\neg \psi(s) \in A) \\ & \vee \exists \psi(x) \in Form_{\mathcal{M}}^1 [\mathcal{M} \models \phi = \forall x \psi] \wedge \forall s \in Tm^c (\psi(s) \in A) \\ & \vee \exists \psi(x) \in Form_{\mathcal{M}}^1 [\mathcal{M} \models \phi = \neg \forall x \psi] \wedge \exists s \in Tm^c (\neg \psi(s) \in A) \end{aligned}$$

Let $\Gamma^M : \mathcal{P}(M) \rightarrow \mathcal{P}(M)$ be the function defined in the following way:

$$\Gamma^M(A) = \{\phi \in \mathcal{M} \mid \Theta_{\mathcal{M}}(\phi, A)\} \quad (\Gamma)$$

As usual, any set $P \subseteq M$ satisfying $\Gamma^M(P) = P$ will be called a *fixpoint* of Γ^M . Such functions generate the extension for the truth predicate of tPT^- in an arbitrary model of PA. More precisely, we have the following well-known fact:

Fact 13. *Let $\mathcal{M} \models PA$. Then*

- (1) *the operator Γ^M is monotone with respect to inclusion, that is, for every sets $A \subseteq B \subseteq M$ we have*

$$\Gamma^M(A) \subseteq \Gamma^M(B),$$

- (2) *there exists a fixpoint of Γ^M ,*
- (3) *if $P \subseteq M$ is any fixpoint of Γ^M , then $(\mathcal{M}, P) \models tPT^-$.*

Our proof of Theorem 10 will be based on the existence of particularly well-behaved fixpoints of Γ in recursively saturated models.

Definition 14. Let $\mathcal{M} \models PA$.

$$\begin{aligned} \Gamma_0^M &= \emptyset \\ \Gamma_{n+1}^M &= \Gamma^M(\Gamma_n^M) \\ \Gamma_\omega^M &= \bigcup_{n \in \omega} \Gamma_n^M \end{aligned}$$

Convention. Instead of $\Gamma^M (\Gamma_n^M, \Gamma_\omega^M)$, we will write $\Gamma (\Gamma_n, \Gamma_\omega)$ omitting the upper index indicating the model.

The following proposition witnesses the first two nice properties of Γ_ω - *if it is a fixpoint of Γ , then it is consistent and satisfies both the extensionality and the normality principles.*¹⁹

Proposition 15. *Let $\mathcal{M} \models PA$ and suppose that Γ_ω is a fixpoint of Γ in M . Then*

$$(\mathcal{M}, \Gamma_\omega) \models RPT^- + \forall \phi \in Sent_{L_{PA}} (\neg(T\phi \wedge T\neg\phi))$$

Proof. Fix $\mathcal{M} \models PA$ and suppose Γ_ω is a fixpoint of Γ . Then by Fact 13

$$(\mathcal{M}, \Gamma_\omega) \models tPT^-$$

That (NORM) holds in $(\mathcal{M}, \Gamma_\omega)$ is straightforward from the definition of Γ . We show consistency first, i.e. we show that for no $\phi \in Sent_{\mathcal{M}}$ it holds that

$$(\mathcal{M}, \Gamma_\omega) \models T(\phi \wedge \neg\phi).$$

¹⁹In fact this proposition is more general: one can show (by a trivial generalisation of our proof) that any least fixpoint of Γ is consistent. Since we do not need it in full generality, we prove the particular case only.

Let us observe that ϕ cannot be an atomic sentence, since PT^- proves that truth is consistent for atomic sentences. Suppose the above is false and take the least $n \in \omega$ such that

$$(\mathcal{M}, \Gamma_n) \models T(\phi \wedge \neg\phi)$$

for some *non-atomic* $\phi \in \text{Sent}_{\mathcal{M}}$. Then $n \neq 0$ since Γ_0 is empty. By considering all possible grammatical forms of ϕ we show that for some $\psi \in \text{Sent}_{\mathcal{M}}$

$$(\mathcal{M}, \Gamma_{n-1}) \models T(\psi \wedge \neg\psi)$$

which contradicts the choice of n .²⁰

Let us now handle (EXT). Suppose for some $t, s \in \text{Term}_{\mathcal{M}}^c$, $\phi(x) \in \text{Sent}_{\mathcal{M}}$ such that $\mathcal{M} \models \text{val}(s) = \text{val}(t)$ we have:

$$(\mathcal{M}, \Gamma_{\omega}) \models T(\phi(s)) \wedge \neg T(\phi(t)).$$

Let n be the least number such that for some formula $\psi(x)$, $\psi(s) \in \Gamma_n$ and $\psi(t) \notin \Gamma_n$. Once again observe that ψ cannot be atomic, since axiom (1) of PT^- guarantees extensionality for such sentences. Moreover $n \neq 0$, since we assumed Γ_n to be nonempty. By considering all the grammatical forms of ψ we show that there is a formula $\theta(x)$ such that

$$(\mathcal{M}, \Gamma_{n-1}) \models T(\theta(s)) \wedge \neg T(\theta(t)).$$

Let us do the step for \vee . If $\psi(x) = \theta_1(x) \vee \theta_2(x)$, then by the assumption concerning $\psi(t)$, for every $k \leq n$:

$$\theta_1(t) \notin \Gamma_k \text{ and } \theta_2(t) \notin \Gamma_k.$$

Since $\psi(s) \in \Gamma_n$, we have that either $\theta_1(s) \in \Gamma_{n-1}$ or $\theta_2(s) \in \Gamma_{n-1}$, which contradicts the choice of n . \square

Let us state our two promised lemmata:

Lemma 16. *If $\mathcal{M} \models \text{PA}$ is recursively saturated, then Γ_{ω} is a fixpoint of Γ .*

Lemma 17. *For every $\mathcal{M} \models \text{PA}$, if Γ_{ω} is a fixpoint of Γ , then $(\mathcal{M}, \Gamma_{\omega}) \models \text{RPT}_{\text{tot}}$.*

Proof of Lemma 16. Let $\mathcal{M} \models \text{PA}$ be recursively saturated. We have to check that $\Gamma(\Gamma_{\omega}) = \Gamma_{\omega}$. Since Γ is monotone it is sufficient to show that

$$\Gamma(\Gamma_{\omega}) \subseteq \Gamma_{\omega}.$$

²⁰For example, if $\phi = \exists x \psi(x)$, then by the definition of Γ_n there must be a number $s \in \mathcal{M}$ such that $(\mathcal{M}, \Gamma_{n-1}) \models T(\psi(s) \wedge \neg\psi(s))$.

Let $\phi \in \Gamma(\Gamma_\omega)$. This means that $\Theta_{\mathcal{M}}(\phi, \Gamma_\omega)$. By considering all the disjuncts constituting $\Theta_{\mathcal{M}}(\phi, \Gamma_\omega)$, we show that $\phi \in \Gamma_\omega$. The only non-trivial steps where we use the recursive saturation of \mathcal{M} are hidden in the cases when $\phi = \forall x\psi$ and $\phi = \neg\exists x\psi$. Let us do the former, the proof of the latter being fully analogous. We have to check that

$$\forall x\psi \in \Gamma_\omega \text{ if and only if for every } t \in Tm_{\mathcal{M}}^c \quad \psi(t) \in \Gamma_\omega. \quad (1)$$

From left to right this is immediate: if $\forall x\psi \in \Gamma_\omega$, then there exists $n \in \omega$ such that $\forall x\psi \in \Gamma_n$. Then, by the definition of Γ it has to be the case that for every $t \in Tm_{\mathcal{M}}^c$

$$\psi(t) \in \Gamma_{n-1}.$$

Since $\Gamma_{n-1} \subseteq \Gamma_\omega$, then the same holds also for Γ_ω (this is also the general pattern of proving the steps for sentential connectives). This ends the proof of the first implication. Suppose now that for every $t \in Tm_{\mathcal{M}}^c$

$$\psi(t) \in \Gamma_\omega. \quad (*)$$

We claim that there is a number $n \in \omega$ such that for every $t \in Tm_{\mathcal{M}}^c$

$$\psi(t) \in \Gamma_n. \quad (**)$$

Aiming at a contradiction, suppose the contrary. Observe that the condition defining Γ_n can be written as an arithmetical formula $\Gamma_n(x)$ such that for every $\phi \in \text{Sent}_{\mathcal{M}}$

$$\phi \in \Gamma_n \text{ if and only if } \mathcal{M} \models \Gamma_n(\phi).^{21}$$

Let us consider the following set:

$$p(x) = \{x \in Tm^c \wedge \neg\Gamma_n(\psi(x)) \mid n \in \omega\}$$

From our assumption that $(**)$ does not hold, together with the fact that for every $k < l \in \omega$:

$$\mathcal{M} \models \forall x(\Gamma_k(x) \rightarrow \Gamma_l(x)),$$

we conclude that $p(x)$ is finitely realisable in \mathcal{M} , so it is a recursive type. By recursive saturation there exists $t \in M$ realising $p(x)$. It follows that $t \in Tm_{\mathcal{M}}^c$ and $\psi(t) \notin \Gamma_\omega$, so we obtain a contradiction with $(*)$.

Now, let $n \in \omega$ be any number satisfying $(**)$. Then by definition of Γ we have

$$\forall x\psi(x) \in \Gamma_{n+1}$$

and therefore

$$\forall x\psi(x) \in \Gamma_\omega,$$

²¹We define $\Gamma_n(x)$ inductively: $\Gamma_0(x) := x \neq x$ and $\Gamma_{n+1}(x) := \Theta(x, \Gamma_n)$, where ‘ $\Theta(x, \Gamma_n)$ ’ is the arithmetical formula obtained from ‘ $\Theta_{\mathcal{M}}(\phi, \Gamma_n)$ ’ by omitting the mention of the model.

which ends the proof. \square

Proof of Lemma 17. Since we know that Γ_ω is a fixpoint of Γ , by Proposition 15 we conclude that

$$(\mathcal{M}, \Gamma_\omega) \models \text{RPT}^-$$

and Γ_ω is consistent. In effect, it is sufficient to show that $(\mathcal{M}, \Gamma_\omega)$ satisfies the internal induction for total formulae. Fix a formula $\phi(x)$ and suppose that

$$(\mathcal{M}, \Gamma_\omega) \models \forall t \in Tm^c(T(\phi(t)) \vee T(\neg\phi(t))).^{22}$$

Then by compositional axioms of PT^- we have $(\mathcal{M}, \Gamma_\omega) \models T(\forall x(\phi(x) \vee \neg\phi(x)))$. Hence there is a number $n \in \omega$ such that

$$\forall x(\phi(x) \vee \neg\phi(x)) \in \Gamma_n.$$

It follows that for every $t \in Tm_{\mathcal{M}}^c$ $\phi(t) \in \Gamma_n \vee \neg\phi(t) \in \Gamma_n$. Since Γ_ω is consistent, we conclude that for every $t \in Tm_{\mathcal{M}}^c$

$$\psi(t) \in \Gamma_n \text{ if and only if } \psi(t) \in \Gamma_\omega.$$

But Γ_n is *arithmetically definable* by formula $\Gamma_n(x)$, so we have the induction axiom for it already in PA. \square

Theorem 10 follows easily from Lemma 16 and Lemma 17.

Remark. We have decided to isolate the two lemmata, because we believe that they both give rise to interesting questions. The first one, based on Lemma 16, is: *what can be said about a model \mathcal{M} given that we know the number of iterations of Γ needed to obtain the least fixpoint?* What we do know is that the converse to Lemma 16 holds, i.e. if Γ_ω is a fixpoint of Γ , then \mathcal{M} is recursively saturated (the proof of this observation is beyond the scope of this paper). The second question, based on Lemma 17, is: *what other fixpoints of Γ give rise to interpretations of PT_{tot} ?* Here we have no clear intuitions.

Let us note one immediate corollary to Theorem 10:

Corollary 18. *CT^- and UTB are not relatively truth definable in PT_{tot} .*

Proof. By theorems of Kaufmann-Schmerl (see [15]) and Smith ([17]), there is a recursively saturated model of PA which does not admit an expansion to a model of CT^- or UTB (a *rather classless* model). By Theorem 10, this model (being recursively saturated) can be expanded to a model of PT_{tot} . Hence, there is a model of PA which expands to a model of PT_{tot} but does not expand to a model of CT^- or UTB . Our corollary follows now from Fact 9. \square

²²Since $(\mathcal{M}, \Gamma_\omega)$ satisfies RPT^- , it makes (EXT) true; therefore not only the compositional axioms for quantifiers, but also the totality condition can be equivalently formulated in its present version, employing the quantification over closed terms.

By Theorem 4, we know that PT_{tot} does not satisfy the semantic conservativity condition. On the other hand, by Theorem 10 we also know that, unlike in the case of CT^- , the interpretation of the truth predicate of PT_{tot} can be found in an arbitrary recursively saturated model of Peano arithmetic. Now, where does it leave Fischer's and Horsten's project?

In view of the aforementioned results, there are two possible moves to consider. One of them consists in rejecting PT_{tot} and searching for another truth theory, satisfying all the demands (a)-(c) from p. 3 of this paper. The second option would involve modifying the demands, in particular, weakening the semantic conservativity requirement.

It is our opinion that the adoption of the second strategy would have far-reaching consequences; namely, that it would amount to nothing less than dropping the original project of providing a characterisation of the use of the truth predicate in model theory. As stressed by Fischer and Horsten, general model theory does not discriminate between models and any description of the notion of model-theoretic truth should take this fact into account. Why then should expandability of all recursively saturated models - but not, say, of prime models - matter for such an endeavour? What is it that permits us to treat model theory as being specifically *about* recursively saturated models? We are not aware of any convincing answer to this question.

Admittedly, expandability of recursively saturated models guarantees the syntactic conservativity of a given theory of truth. Indeed, syntactic conservativity is crucial for a related but different philosophical project, proposed by Fischer in [5] and [7], where the core idea is to present truth as an instrumental device, on a par with 'ideal elements in mathematics' in Hilbert's programme (cf. [7], p. 294). In such a context there is still a place for theories like PT_{tot} . Nevertheless, we should emphasise that the present project - that of characterising the use of the truth predicate in model theory - is different and we just do not see how it could proceed without the semantic conservativity condition.

In effect, from the philosophical point of view the first option definitely seems more attractive. Moreover, some axiomatic truth theories in the vicinity of PT_{tot} merit further attention as very promising candidates for the role of a theory realising Fischer's and Horsten's postulates.²³

Acknowledgments. The research presented in this paper was supported by the National Science Centre, Poland (NCN), grant number 2014/13/B/HS1/02892.

²³One of these promising theories is WPT_{tot} , which is obtained from PT_{tot} by modifying the compositional truth axioms in accordance with the principles of Weak Kleene logic. However, the analysis of the properties of WPT_{tot} and related theories goes beyond the scope of the present paper.

References

- [1] Cantini, A. (1989), “Notes on formal theories of truth”, *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik*, vol. 35(2), 97–130.
- [2] Cieśliński, C. (2015), “Typed and untyped disquotational truth”, in T. Achourioti, H. Galinon, K. Fujimoto, and J. Martínez-Fernández (eds), *Unifying the Philosophy of Truth*, 307–320. Springer.
- [3] Cieśliński, C. (2015), “The innocence of truth”, *dialectica*, vol. 69(1), 61–85.
- [4] Cieśliński, C. (2011), “T-equivalences for positive sentences”, *The Review of Symbolic Logic*, vol. 4(2), 319–325.
- [5] Fischer, M. (2014), “Truth and speed-up”, *The Review of Symbolic Logic*, vol. 7(2), 319–340.
- [6] Fischer, M. (2009), “Minimal truth and interpretability”, *The Review of Symbolic Logic*, vol. 2(04), 799–815.
- [7] Fischer, M. (2015) “Deflationism and instrumentalism”, in T. Achourioti, H. Galinon, K. Fujimoto and J. Martínez-Fernández (eds.) *Unifying the Philosophy of Truth*, Springer, 293–306.
- [8] Fischer, M. and Horsten, L. (2015), “The expressive power of truth”, *The Review of Symbolic Logic*, vol. 8(2), 345–369.
- [9] Fujimoto, K. (2010), “Relative truth definability of axiomatic truth theories”, *The Bulletin of Symbolic Logic*, vol. 16, 305–344.
- [10] Halbach, V. (2009), “Reducing compositional to disquotational truth”, *The Review of Symbolic Logic*, vol. 2(4), 786–798.
- [11] Halbach, V. (2011), *Axiomatic Theories of Truth*, Cambridge University Press.
- [12] Horsten, L. (1995), “The semantical paradoxes, the neutrality of truth and the neutrality of the minimalist theory of truth”, in P. Cartois (ed.), *The Many Problems of Realism*, vol. 3 of *Studies in the General Philosophy of Science*, Tilburg University Press, 173–187.
- [13] Kaye, R. (1991), *Models of Peano Arithmetic*, Clarendon Press.
- [14] Ketland, J. (1999), “Deflationism and Tarski’s paradise”, *Mind*, vol. 108(429), 69–94.
- [15] Kossak, R., Schmerl, J. H. (2006), *The Structure of Models of Peano Arithmetic*, Clarendon Press, Oxford.

- [16] Shapiro, S. (1998), “Proof and truth: through thick and thin”, *The Journal of Philosophy*, vol. 95(10), 493–521.
- [17] Smith, S. T. (1989), “Nonstandard definability”, *Annals of Pure and Applied Logic*, vol. 42, 21–43.